

Comment to "General Solution to Unidimensional Hamilton-Jacobi Equation" (arXiv:1302.0591v1)

Irina YEHOCHENKO

Institute of Mathematics of NAS Ukraine, 3 Tereshchenkivs'ka Str.,
01601 Kyiv-4, Ukraine E-mail: iyegorch@imath.kiev.ua

Abstract

We present previous results on the general solution of the multidimensional Hamilton-Jacobi equation $\frac{\partial u}{\partial t} - \frac{\partial u}{\partial x_a} \frac{\partial u}{\partial x_a} = 0$ and methods that were used to find such general solution.

1 Introduction

The paper [1] says that *"Unfortunately, due to the nonlinearity of this equation (the Hamilton Jacobi equation - IY) till now there is no available technique to determine a general solution"*.

I would like to point out that such technique does exist (the method of the hodograph transformation), and it was used by us in 1991 [2] (following [3]) to find the general solution for the Hamilton-Jacobi equation

$$\frac{\partial u}{\partial t} - \frac{\partial u}{\partial x_a} \frac{\partial u}{\partial x_a} = 0. \quad (1)$$

Here $a = 1, \dots, n$ (so, we deal with the arbitrary number of the space variables).

2 General solution of the Hamilton-Jacobi equation

Finding of the general solution of the Hamilton-Jacobi equation in [2] was regarded as an intermediate task and was not specifically acknowledged, so it might have been overlooked. However, it is a technical problem, and I really could not claim that our paper [2] was the first instance when such

general solution was mentioned. General solution of the Hamilton-Jacobi equation in a different form was considered in [4].

Anyway, such general solution of the rank n (that means that $\det\{\frac{\partial u^2}{\partial x_a \partial x_b}\} \neq 0$) has the form

$$u = x_a y_a - \frac{t}{2} y_a y_a + \Phi(\vec{y}), \quad (2)$$

$$x_a - t y_a + \frac{\partial \Phi}{\partial y_a} = 0, \quad (3)$$

$y = (y_1, \dots, y_n)$ are parameter functions, Φ is an arbitrary function of its parameters subject to the condition (3). We assume summation from 1 to n by the repeated indices.

Rank of the solution is the rank of the matrix $\{\frac{\partial u^2}{\partial x_a \partial x_b}\}$.

3 Hodograph transformation and its use for finding of solutions of PDE

The method of the hodograph transformation (interchanging of some of the dependent and independent variables) in the equation was described e.g. in [5].

The hodograph transformations for the equation (1) and $\text{rank}\{\frac{\partial u^2}{\partial x_a \partial x_b}\} = n$ is as follows:

$$y_0 = t, y_a = \frac{\partial u}{\partial x_a}, \quad (4)$$

$$H(y) = x_a \frac{\partial u}{\partial x_a} - u,$$

Then we come to the Hamilton-Jacobi equation in new variables

$$\frac{\partial H}{\partial y_0} = -y_a y_a.$$

The general solution in terms of the function H :

$$H = -ty_a y_a - \Phi(\overrightarrow{y}). \quad (5)$$

Using to the expression (5) the inverse hodograph transformations (4), we will receive the solution (2) with the conditions (3).

We considered the case of solutions of the rank n ; the procedure for the ranks from 0 to $n - 1$ is similar with the relevant number of the parameter functions.

The hodograph transformations were used in [3] for finding of the general solution of the system of eikonal and d'Alembert equations.

As to utilisation of the hodograph transformations for solving of other PDEs and systems of PDEs, see [6], [7].

The method presented in [1] is technically the same as the hodograph transformation, but in different terminology, and gives the same result in different notations formula (5) in [1]).

The hodograph transformations is a straightforward and powerful technique to linearise and solve nonlinear PDEs (not only first-order), and probably worth writing of a detailed tutorial.

References

- [1] Lewtchuk Espindola Maria, General Solution to Unidimensional Hamilton-Jacobi Equation. arXiv:1302.0591v1
- [2] Fushchych W.I., Yehorchenko I.A., Non-Lie ansätze and conditional symmetry of the nonlinear Schrödinger equation, Ukrainian Math. Journal, 1991, vol. 43, no. 12, p.1507-1515.
- [3] Fushchych W.I., Zhdanov R.Z. and Revenko I.V., General solutions of the nonlinear wave and eikonal equations, Ukrainian Math.J., 1991, vol. 43, no.11, p. 1364-1379
- [4] Epstein S.T., General Solutions of the Hamilton-Jacobi Equation, Amer.J.Phys., 1964, Vol. 32, p. 688-690.
- [5] Courant R., Friedrichs K. O., Supersonic Flow and Shock Waves, Springer, 1948.
- [6] Meleshko S. V., Methods for Constructing Exact Solutions of Partial Differential Equations: Mathematical and Analytical Techniques with Applications to Engineering, Springer, 2005.
- [7] Kevorkian J., Partial Differential Equations: Analytical Solution Techniques (Texts in Applied Mathematics), Springer, 1999.